

FOREVER CHEMICALS:

HOW WE ARE TACKLING
HARMFUL EXPOSURE -
AT A GLANCE



1,300 –
mother-
child pairs
we are
following
in HELIX
project

80,000 – mother-
child pairs
we are
following in
ATHLETE
study



11+ – European countries
collaborating




250,000 – children and
parents in LifeCycle project



89 – outdoor, chemical,
and lifestyle exposures
measured during
pregnancy



3 – major recommendations
for policymakers and **3** for
the general public



EVIDENCE BRIEFING

HARMFUL EXPOSURE: CHEMICALS AND HEALTH



KEY CONTACT

Tiffany Yang

tiffany.yang@bthft.nhs.uk

WHAT IS THE PROBLEM?

Chemicals are all around us in the form of the **objects we touch, the air we breathe, the products we use, and the food and liquids we consume**.

Some chemicals are of concern because of the **negative health impacts** they have, such as lead, arsenic, and other naturally-occurring heavy metals on cognitive function and behaviour^{1,2}.

These are **especially concerning** when exposures occur at sensitive periods of development, such as during **pregnancy** or **in childhood**.

The increasing use of synthetic chemicals such as phthalates and phenols in industrial and everyday consumer products to act as solvents, fixatives, preservatives, and plasticisers has also become a **cause for concern**^{3,4}.

These chemicals are found in our **personal care products** (shampoos, soaps, lotions/creams, perfumes, etc.) as well as **common household objects** (vinyl flooring, plastic food storage containers and packaging, etc.).

A growing body of evidence suggests that some of these compounds act as endocrine-disrupting hormones, meaning they have the **potential to disrupt** common hormone pathways such as the thyroid, androgen, and oestrogen pathways⁴⁻⁸.

Exposure to these chemicals have been associated with a range of health implications including **obesity, impaired glucose tolerance, gestational diabetes, and semen quality**⁹.





WHAT HAVE WE FOUND?

Born in Bradford has been involved in **several European research collaborations** which have looked at the links between **exposures to chemicals from our environment** and **health**.

These collaborations have included the **HELIX (Human Early Life Exposome) study**, a collaboration between **six birth cohorts in six European countries** (UK, Spain, Norway, France, Lithuania, Greece) which followed **1,300 mother-child pairs**¹⁰.

We also played a key role in its follow-up and expansion to **16 existing cohorts in 11 European countries** with **80,000 mother-child pairs** in the **ATHLETE (Advancing Tools for Human Early Lifecourse Exposome Research and Translation)** study¹¹.

We have also been a partner in the **LifeCycle project**, which established the EU Child Cohort Network of harmonised data from **19 pregnancy and childhood cohorts** with **more than 250,000 children and their parents**¹².

Through these studies, newly-collected data using biological samples such as **urines and blood**, as well as **questionnaire data**, has allowed epidemiological analyses to **answer questions** about the amount of exposure experienced by

BiB families and the impact of these exposures on health.

We are routinely exposed to a wide range of environmental chemicals from foetal life through adulthood

We are exposed to a wide range of both **persistent** and **non-persistent chemicals**¹³.

Some chemicals, like per- and polyfluoralkyl substances (PFAS), which are used in **non-stick cookware, packaging**, and make **fabrics stain-resistant**, are known as **“forever chemicals”** because they **persist in the environment** or **your body** long after they are used or consumed¹⁴.

Other chemicals, such as Bisphenol A or phthalates, which are used in **plastic containers, medical equipment**, and **personal care products**, don't persist in the environment and have short half-lives in the body, but due to **almost constant exposure** we retain a **body burden** of these chemicals^{15,16}.

The types of exposures differed between **countries** as well as by **age group**.

Some of this difference can be explained by **different sources of exposure for adults compared to children**,

such as the use of personal care products and cosmetics among adults, or, among children, because of greater inhalation, dermal contact, or dietary intake relative to body size¹⁷.

Exposure to heavy metals and chemicals can negatively impact on behaviour, blood pressure, respiratory health, and liver function in children

We know that foetal life and childhood are important periods for health as they are periods of **rapid development** and adverse events during these periods could have **long-term impacts on health**.

BiB was part of a **novel study** that was the first to look simultaneously at a range of **89 outdoor, chemical, and lifestyle exposures during pregnancy** and **128 exposures during childhood** on different health outcomes in childhood:

- We found that exposure to both persistent (PFAS) and non-persistent (BPA) chemicals and metals (copper) are associated with **increased blood pressure** in children whereas persistent organochlorine compounds such as PCBs, as well as non-persistent compounds like phthalates, have been **inversely associated with childhood blood pressure**¹⁸.
- We found that exposure to BPA, phthalates, and DETP, a pesticide known to be neurotoxic, in the foetal period was associated with **worse externalising behaviour in childhood**¹⁹. Heavy metals such as **lead** and **copper** were also associated with worse childhood behaviour such as scoring higher on an **ADHD index**²⁰.

- We found that BPA exposure in the foetal period was associated with higher odds of **childhood wheeze** and **asthma**²¹. Worse **childhood lung function** was also linked with exposures to PFAS, copper, parabens (a common preservative used in personal care products, pharmaceuticals, and the food industry), and phthalates²². **Exposure to copper** has also been implicated in **childhood obesity**²³.
- We found that exposure to PFAS in the foetal period was linked to an increased risk of **liver injury in childhood**²⁴.

Some chemicals found in foods, or formed when foods are cooked, can have harmful effects for our health

- Acrylamide is a neurotoxin and probable **cancer-causing** agent; exposure to this compound has been linked to intakes of starchy foods that have been fried or baked at high temperatures. In the BiB cohort, **chips, toast, and crisps** were the greatest contributors to intakes of acrylamide, with **White British women** more likely to have higher intakes with increasing levels of **deprivation**²⁵. Dietary intakes of acrylamide was associated with a **reduced birth weight** and **smaller head circumference**²⁶.



- **Fish intake** is beneficial for health but excessive intake of fish (more than four times every week) can increase the amount of exposure to persistent chemicals and heavy metals found in the fish, such as polychlorinated bisphenyls (PCBs), arsenic, and mercury compared to the recommended 2-3 times/week²⁷. By keeping to recommended intakes of fish, rather than low or excessive intakes, a **better metabolic profile was observed in children**, with a decrease in a metabolic syndrome score as well as lower levels of inflammatory biomarkers²⁸.
- **Pesticides** are commonly used in agriculture and we are exposed to these pesticides through **food consumption**. **Fruit intake** increases exposures to organophosphate pesticides but consuming **organic foods** reduces these exposures.

Individuals can lower their exposure to some chemicals by swapping out plastic food storage containers and choosing alternative personal care products

We are constantly exposed to chemicals in **our daily life**. Some of these chemicals, like BPA, parabens, and phthalates, don't stay in the body for long periods of time after exposure but because they are used in everyday products like **cosmetics, lotions, creams, toothpastes, shampoos, food containers and food packaging**, we are **repeatedly exposing ourselves** to these chemicals. We undertook a review of the current scientific literature to understand

whether it is possible to reduce our exposure to these chemicals through actions that we can take.

We found that it is possible to reduce our exposures by **replacing plastic food and beverage containers with glass or stainless steel, looking at the labels on our cosmetics and personal care products to see if they contain the chemicals**, or **decreasing intakes of canned and packaged foods** ^[29].

WHAT HAS CHANGED AS A RESULT?

Our findings have helped **increase our understanding** of how our **everyday exposures can impact on our health and the health of our children**.

This has **prompted a human study** to understand whether feasible and effective interventions can be created to **reduce our exposures to chemicals** found in our **cosmetics and personal care products**.

Our work has also contributed to national, international, and US changes in guidelines to reduce exposures to acrylamide.



RECOMMENDATIONS FOR POLICYMAKERS BASED ON OUR EVIDENCE

- That **consumer information on the presence of chemicals** used within products and packaging are **made accessible** and **widely available**
- That policy is implemented to **target the use of chemicals** in multiple sectors including **processing, manufacturing, and packaging** to reduce individual burden in identifying and finding alternative products
- That **greater consideration to the potential harms of chemicals**, such as **pesticides** and **preservatives**, is given prior to these chemicals being introduced into manufacturing processes or consumer products

ADDITIONAL RECOMMENDATIONS FOR THE GENERAL PUBLIC

- Try to use **fewer plastic food and beverage storage containers**, especially when **microwaving**, and switch to non-reactive materials like glass or stainless steel if you can
- Try to **reduce your consumption of tinned foods and foods stored in plastic packaging** because chemicals in packaging can migrate into the food
- Keep to the **recommended fish intake of 2-3 times/week** and switch to **organic foods** if you can
- When buying personal care products, look at the ingredients list and try to **avoid products with triclosan, phthalates, and parabens** which are known to impact health



REFERENCES

1. Fu, Z. & Xi, S. The effects of heavy metals on human metabolism. *Toxicol. Mech. Methods* **30**, 167–176 (2020).
2. Al osman, M., Yang, F. & Massey, I. Y. Exposure routes and health effects of heavy metals on children. *BioMetals* **32**, 563–573 (2019).
3. Dodson, R. E. *et al.* Endocrine disruptors and asthma-associated chemicals in consumer products. *Environ. Health Perspect.* **120**, 935–43 (2012).
4. Gore, A. C. *et al.* EDC-2: The Endocrine Society's Second Scientific Statement on Endocrine-Disrupting Chemicals. *Endocr. Rev.* **36**, E1–E150 (2015).
5. Kahn, L. G., Philippat, C., Nakayama, S. F., Slama, R. & Trasande, L. Endocrine-disrupting chemicals: implications for human health. *Lancet Diabetes Endocrinol.* **8**, 703–718 (2020).
6. Meneguzzi, A., Fava, C., Castelli, M. & Minuz, P. Exposure to Perfluoroalkyl Chemicals and Cardiovascular Disease: Experimental and Epidemiological Evidence. *Front. Endocrinol. (Lausanne)*. **12**, 1–13 (2021).
7. Vandenberg, L. N. *et al.* Hormones and endocrine-disrupting chemicals: low-dose effects and nonmonotonic dose responses. *Endocr. Rev.* **33**, 378–455 (2012).
8. Ding, D. *et al.* The EDKB: an established knowledge base for endocrine disrupting chemicals. *BMC Bioinformatics* **11**, 5–11 (2010).
9. Walker, L. E. The importance of environmental embodiment for public health professionals: Stress triggers, environmental toxicants, and strategies for education. *Diss. Abstr. Int. Sect. B Sci. Eng.* **77**, No-Specified (2016).
10. Maitre, L. *et al.* Human Early Life Exposome (HELIX) study: a European population-based exposome cohort. *BMJ Open* **8**, e021311 (2018).
11. Vrijheid, M. *et al.* Advancing tools for human early lifecourse exposome research and translation (ATHLETE). *Environ. Epidemiol.* **5**, E166 (2021).
12. Jaddoe, V. W. V. *et al.* The LifeCycle Project-EU Child Cohort Network: a federated analysis infrastructure and harmonized data of more than 250,000 children and parents. *Eur. J. Epidemiol.* **35**, 709–724 (2020).
13. Haug, L. S. *et al.* In-utero and childhood chemical exposome in six European mother-child cohorts. *Environ. Int.* **121**, 751–763 (2018).
14. CHEM Trust. PFAS – the 'forever chemicals', Invisible threats from persistent chemicals. A CHEM Trust briefing [DEG3]. 1–18 (2019).
15. Völkel, W., Colnot, T., Csanády, G. A., Filser, J. G. & Dekant, W. Metabolism and kinetics of bisphenol a in humans at low doses following oral administration. *Chem. Res. Toxicol.* **15**, 1281–7 (2002).
16. Koch, H. M., Bolt, H. M., Preuss, R. & Angerer, J. New metabolites of di(2-ethylhexyl)phthalate (DEHP) in human urine and serum after single oral doses of deuterium-labelled DEHP. *Arch. Toxicol.* **79**, 367–76 (2005).
17. Snodgrass, W. Physiological and biochemical differences between children and adults as determinants of toxic exposure to environmental pollutants. in *Similarities and Differences between children and adults: implications for risk assessment* (eds. P, G., C, H. & S, O.) 35–42 (International Life Sciences Institute, 1992).
18. Warembourg, C. *et al.* Early-Life Environmental Exposures and Blood Pressure in Children. *J. Am. Coll. Cardiol.* **74**, 1317–1328 (2019).
19. Jedynak, P. *et al.* Prenatal exposure to a wide range of environmental chemicals and child behaviour between 3 and 7 years of age – An exposome-based approach in 5 European cohorts. *Sci. Total Environ.* **763**, (2021).
20. Maitre, L. *et al.* Early-life environmental exposure determinants of child behavior in Europe: A longitudinal, population-based study. *Environ. Int.* **153**, (2021).
21. Abellan, A. *et al.* In utero exposure to bisphenols and asthma, wheeze, and lung function in school-age children: a prospective meta-analysis of 8 European birth cohorts. *Environ. Int.* **162**, (2022).
22. Agier, L. *et al.* Early-life exposome and lung function in children in Europe: an analysis of data from the longitudinal, population-based HELIX cohort. *Lancet Planet. Heal.* **3**, e81–e92 (2019).
23. Vrijheid, M. *et al.* Early-life environmental exposures and childhood obesity: An exposome-wide approach. *Environ. Health Perspect.* **128**, 1–14 (2020).
24. Stratakis, N. *et al.* Prenatal Exposure to Perfluoroalkyl Substances Associated With Increased Susceptibility to Liver Injury in Children. *Hepatology* **72**, 1758–1770 (2020).
25. Hepworth, S. J. *et al.* P20 Description of dietary intakes of acrylamide in the born in Bradford birth cohort study. *J. Epidemiol. Community Heal.* **64**, A41–A41 (2010).
26. Pedersen, M. *et al.* Birth weight, head circumference, and prenatal exposure to acrylamide from maternal diet: The European prospective mother-child study (NewGeneris). *Environ. Health Perspect.* **120**, 1739–1745 (2012).
27. Papadopoulou, E. *et al.* Diet as a source of exposure to environmental contaminants for pregnant women and children from Six European Countries. *Environ. Health Perspect.* **127**, 1–13 (2019).
28. Stratakis, N. *et al.* Association of Fish Consumption and Mercury Exposure During Pregnancy With Metabolic Health and Inflammatory Biomarkers in Children. *JAMA Netw. open* **3**, e201007 (2020).
29. <https://pubmed.ncbi.nlm.nih.gov/36988899/>